

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.

THIS PAGE BLANK (USP 10)

18



Europäisches Patentamt
European Patent Office
Office européen des brevets

11 Publication number:

0 146 742
A1

12

EUROPEAN PATENT APPLICATION

21 Application number: 84113349.9

51 Int. Cl.⁴: A 61 K 31/12
A 61 K 47/00

22 Date of filing: 06.11.84

30 Priority: 08.11.83 JP 208395/83

43 Date of publication of application:
03.07.85 Bulletin 85/27

64 Designated Contracting States:
BE CH DE FR GB IT LI NL SE

71 Applicant: Eisai Co., Ltd.
6-10, Kojishikawa 4-chome Bunkyo-ku
Tokyo 112(JP)

72 Inventor: Okuyama, Shinichi
15-38, Nakayama 3-chome
Sensai-shi Miyagi Prefecture(JP)

72 Inventor: Mishina, Hitoshi
7-5, Aramaki-chuo
Sendai-shi Miyagi Prefecture(JP)

72 Inventor: Furuse, Kazumaro
3-5, Inogashira 2-chome
Mitaka-shi Tokyo(JP)

72 Inventor: Murotani, Isao
5-11, Shodo 2-chome Totsuka-ku
Yokohama-shi Kanagawa Prefecture(JP)

74 Representative: Hansen, Bernd, Dr.rer.nat. et al,
Hoffmann, Eitle & Partner Patentanwälte Arabellastrasse
4
D-8000 München 81(DE)

64 Therapeutic preparation containing ubidecarenone.

67 Cutaneous administration of ubidecarenone, for example, application of ubidecarenone in the form of an ointment has been proven to be effective for treatment of various skin disorders caused by radiation, such as radiation ulcers and radiation dermatitis.

0146742

THERAPEUTIC PREPARATION CONTAINING UBIDECARENONE

- 1 -

This invention relates to a therapeutic preparation
5 for radiation ulcers which contains ubidecarenone as an
effective component and also to a therapeutic method for
radiation ulcers which features use of ubidecarenone.

The radiotherapy for cancers has conventionally been
10 prone to pay principal attention to the killing of cancer
cells and to disregard the need for treatment of secondary
problems or troubles developed on the skin by radiation,
namely, radiation ulcers, radiation dermatitis or the like.

Generally speaking, skins which have been subjected
15 to the radiotherapy are accompanied by intensive atrophy and
their vascular walls have been rendered weak. These skins
tend to undergo damages even by slightest extrinsic stimuli.
The radiotherapy brings about such inherent side effects
that it induces local obstructive arterioloarteritis,
20 destroys lymph vessels and the like and decreases
SH-containing enzymes at certain local sites, thereby making
the circulation of body fluid poor. On the other hand,
strong tissues such as nervous tissues are able to survive
the radiotherapy. Therefore, a metabolic disorder developed
25 by poor circulation at a local site does not immediately
stimulate nerves. If a disorder should be developed by an
external cause such as extrinsic force on a skin which has
been subjected to the radiotherapy, the disorder will be

0146742

intractable and moreover, will spread to its surrounding skin. In addition, the patient will suffer from severe irritation and pains and the danger of mixed infection will increase as time goes on.

5 It is believed to be essential to achieve an immediate improvement to the circulation at such an infected site in order to treat the above-mentioned state of radiation ulcer or radiation dermatitis. Since such an infected site is clotted by inflammation products and the like, its treatment has hitherto been carried out by applying either singly or in combination the hyperbaric oxygen therapy, the glutathione injection therapy, the ascorbic acid and urokinase injection therapy, the oral cytochrome c administration therapy and so on. Furthermore, 15 patients of this sort often develop some disorders at their hearts, livers, etc. Upon selection of drugs for radiation ulcers or radiation dermatitis, it is thus important to pay attention so that such additional disorders will also be cured without development of side effects. It is also 20 recommended to prevent recurrence of such problems by using a tissue activator or the like even after the curing of radiation ulcers or radiation dermatitis, because the activity of the tissue of such an infected skin has been lowered.

25 The following publications 1) - 7), all in Japanese except for 5) and 7), will be listed by way of example for

furth r-detailed explanation of the above prior art findings.

0146742

- 1) Shiojima, S., Mishina H. and Ohuchi I.: Present Situation of Hyperbaric Oxygen Therapy in Tohoku Rosai Hospital. Nichiroshi, 21(8), 359-365, (1973);
- 2) Mishina H., Haryu T., Shiojima S. and Imaizumi A.: Clinical Example of Radiation Ulcers. Shinryo to Shinyaku, 11(1), 105-110, (1974);
- 3) Abe, Y., Shima T., Akiyama K. and Ohga H.: Treatment of Radiation Skin Disorders. Rinpo, 2(12): 19-25, (1967);
- 4) Mishina H. and Haryu T.: Results of Application of Urokinase for Postoperative Breast Cancer after ⁶⁰Co Radiation. Medical View 7(1), 25-26, (1972);
- 5) Guettier, Y. et al: Cahiers D'O.R.L., 1, 77, (1966);
- 6) Mishina H., Haryu T., Shiojima S., Imaizumi A. and Sato, T.: Hyperbaric Oxygen Therapy of Roentgen Ulcers. Basic Pharmacol Therapeut. 5, 1439, (1977); and
- 7) Sapiro, B.: Biochemical Mechanism in the Action of Radiation. In The Biological Basis of Radiation Therapy, E. E. Schwartz, ed., Lippincott, Philadelphia, 1966, pp. 31-59.

0146742

As mentioned above, radiation disorders are unavoidable for the carcinostatic radiation therapy. These disorders are intractable especially where they are accompanied by loss of tissue. Where scar tissue is spread over a wide area for example in a radiation ulcer, the epithelialization has proceeded to a considerable degree and the thus-infected skin has become weak against physical, chemical and biological influence. Therefore, its curing has become more difficult.

10

With the foregoing circumstances in view, the present inventors have studied various methods, which were led by the above-mentioned prior art methods, with a view toward achieving the treatment of radiation ulcers. As a result, it has been found that an administration of ubidecarenone to an ulcerated site leads to excellent curing results, leading to completion of this invention.

Accordingly, the overall object of this invention is to cure radiation ulcers. More specifically, an object of this invention is to provide therapeutic preparation and method effective for the curing of radiation ulcers.

In one aspect of this invention, there is thus provided a cutaneously-applicable therapeutic preparation for radiation ulcers which contains ubidecarenone as an effective component.

0146742

In another aspect of this invention, there is also provided a therapeutic method for radiation ulcers, which comprises cutaneously administering ubidecarenone.

5 The therapeutic preparation and method of this invention are effective for treating and curing radiation ulcers without development of noticeable side effects.

The above and other objects, features and advantages of the present invention will become apparent from the following description and the appended claims.

10

The term "radiation ulcers" as used herein should be interpreted in broad sense. It means all disorders which skins have developed at radiated sites. Therefore, the
15 above term embraces not only those proceeded to actual ulcers but also those including radiation dermatitis. In other words, the effectiveness of the present invention has been proven as a result of an observation on its effects which observation was carried out using as samples those
20 already proceeded to ulcers. As readily envisaged from the above observation, the present invention is also effective for intermediary disorders in each of which the destruction of tissue has not proceeded to an ulcer.

Ubidecarenone is also called ubiquinontene or
25 coenzyme Q₁₀ and has already been used as a therapeutic agent for congestive heart failure in the pharmaceutical field. The same ubidecarenone is also useful in the

0146742

practice of this invention. Ubidecarenone is extracted from mitochondria of bovine cardiac muscle and is known to take part in the electron transport system. Corollary to this, ubidecarenone can improve the percentage oxygen utilization of cardiac muscle and maintain the ATP-yielding function at a high level even when the cardiac muscle is in an ischemic state. As a result, it has been known that ubidecarenone can reduce disorders, which the tissue of ischemic cardiac muscle would develop, and can hence improve the drop of heart contraction function.

There has however been absolutely unknown the fact that radiation ulcers can be successfully cured by cutaneously-administered ubidecarenone. This surprising effects of ubidecarenone have been uncovered for the first time by the present inventors.

Ubidecarenone is crystalline powder of a yellow to orange color with a melting point of 48 - 52°C. It is oil-soluble but is hardly soluble in water or methanol. As mentioned above, it has hitherto been perorally administered for the improvement of various symptom of congestive heart failure. For the sake of reference, its subacute toxicity and chronic toxicity upon peroral administration are as follows:

Subacute toxicity:

Ubidecarenone was perorally administered at 40, 200 and 1,000 mg/kg/day to male and female Wistar rats continuously over a period of 5 weeks and also at 60 and 600

0146742

mg/kg/day to male and female rabbits continuously over a period of 23 days. No differences were observed in general appearance, blood, urine test and morphological observation (both visual and histological) between the thus-treated rats and rabbits and those of a control.

Chronic toxicity:

Ubidecarenone was forcedly and perorally administered at 6, 60 and 600 mg/kg/day to female and male Wistar rats continuously over a straight period of 26 weeks. As a result, no differences were observed in general appearance, blood, urine test and morphological observation (both visual and histological) between the thus-treated rats and those of a control.

The present invention features cutaneous administration of ubidecarenone to a site where such a radiation ulcer as defined above has developed.

For its cutaneous administration, ubidecarenone may be applied, as is, directly to such an infected site but it is desirable to apply it in the form of a preparation suitable for its coating onto the skin whenever possible. Furthermore, it is also feasible to administer ubidecarenone in combination with other drugs, for example, cytochrome c, urokinase and/or the like. It should however be borne in mind that the present invention is not necessarily limited to such a combined use of other drugs.

The content of ubidecarenone in the therapeutic preparation according to this invention may be recommended

0146742

to be 0.05 - 5.0% or more preferably, 0.1 - 2.0%. The therapeutic preparation may be applied in a suitable amount, depending on the size and progress of each ulcer.

5 When applied to the skin, ubidecarenone exhibits good stability and little irritation to the skin. Table 1 shows by way of example results of various tests, i.e., the primary skin irritation, cumulative irritation, blepharo-irritation, photo-toxicity, challenge, optical challenge and patch test.

10

0146742

Test	Conc.	Solvent	Results/Conclusion	
Primary skin irritation	1%	Squalane	0.1	Little skin irritation
	-	ditto	0.1	ditto
Cumulative skin irritation	1%	ditto	0.3	ditto
	-	ditto	0.3	ditto
Blepharo-irritation	1%	ditto	Little blepharo-irritation	
	-	ditto	ditto	
Phototoxicity	1%	ditto	(-)	Little phototoxicity
	10%	ditto	(-)	ditto
Challenge (Adjuvante & Patch Method)	Induction: 5% Acetone			
	5%	Acetone	0/10	Little challenge
	1%	ditto	0/10	Little challenge
Photo-challenge (Adjuvante Strip method)	Induction: 10% Acetone			
	10%	Acetone	0/5	Little photo-challenge
	5	ditto	0/5	ditto
	2	ditto	0/5	ditto
	1	ditto	0/5	ditto
	0.5	ditto	0/5	ditto
Patch test	1%	Squalane	0/54	Little skin irritation
	-	ditto	0/54	ditto

0146742

In order to form ubidecarenone into a preparation suitable for cutaneous administration, one or more excipients or vehicles which causes little irritation may be suitably chosen. For example, glycerin, hohoba oil, cetyl
5 alcohol, olive oil, a glyceryl fatty acid ester and/or the like may be selected to prepare a preparation for cutaneous administration in a manner known per se in the art.

Effects of the present invention will next be described in the following case reports:

10 Case 1:

Female, 68 y.o.a. The patient underwent a surgery on heft left breast cancer. It recurred in her left chest wall seven years later, resulting in the formation of a carcinomatous ulcer. This carcinomatous ulcer became
15 resistant to the cryosurgery and carcinostatic chemotherapy, leading to need for the radiotherapy. The patient was exposed to a total radiation dose of 40 Gy while using oily bleomycin and carmofur ointment in combination with a view toward enhancing the effects of the radiotherapy. Upon
20 peroral administration of zinc sulfate and injection of intaserine to facilitate wound healing, the ulcer became cancer-free and an eschar was produced. However, she fell down accidentally at this stage and the wound was traumatized and ulcerated again. Thus, cythochrome c was injected and at
25 the same time, Bendazac ointment was applied. The epithelialization did not proceed any further although some erythema and hyperemia were still observed at the infected

0146742

site. At this stage, the coating of a therapeutic preparation to be described in Example 1, which pertains to the present invention, was started. Effects of the treatment were observed within 24 hours as reduction to spontaneous pains and exudate. Four days later, the epithelialization of the surrounding area was observed together with the formation of benign granulation tissue over the surface of the ulcer. The wound was covered by a thick cuticle before the 20th day, resulting in disappearance of the ulcer. The thus-cured skin was thereafter resistant to usual extrinsic forces.

Case 2:

Female, 73 y.o.a. This patient was subjected to postoperative radiation for the treatment of cervical cancer 18 years before. Her lower abdomen was infected by a radiation ulcer. Some scars were formed there. It was atrophic. Besides pigmentation, telangiectasia and ulcer formation were also observed. In the Case, the hyperbaric oxygen therapy, zinc sulfate, injectable glutathione, injectable intaserine and Bendazac ointment were applied intermittently in combination. However, the ulcer was not improved. The oral ubidecarenone therapy as also applied for further 2 months but no change was observed as to the ulcer. Thus, the coating of a therapeutic preparation to be described in Example 1, which pertains to this invention, was initiated. The ulcer healed up in two weeks and the pigmentation and telangietasia were improved.

0146742

The invention will hereinafter be described more specifically in the following Examples:

Example 1:

	Stearyl alcohol	5 wt. %
5	Stearic acid	2
	Hydrogenated lanolin	2
	Squalane	6
	Isopropyl myristate	4
	Polyoxyethylene (25 moles) cetyl alcohol ether	3
10	Glycerin monostearate	2
	Ubidecarenone	0.3
	Propylene glycol	5
	Butyl paraben	0.2
15	Antioxidant	as needed
	Perfume	as needed
	Purified water	70.5

The above components were mixed in a manner known per se in the art into a homogeneous cream as a
20 therapeutic preparation of this invention.

Example 2:

	Cetyl alcohol	4 wt. %
	Behenic acid	2.5
	Vaseline	3
25	Liquid paraffin	10
	2-Octyldodecyl alcohol	4
	Bees wax	1

0146742

	Glycerin monstearate	2
	Polyoxyethylene(20 moles) sorbitan monolaurate	2
	Ubidecarenone	1
5	Glycerin	5
	Polyethyelene glycol 1500	5
	Caustic potash	0.2
	Ethylparaben	0.4
	Perfume	as needed
0	Purified water	59.9

The above components were mixed in a manner known
per se in the art into a homogeneous cream as a
therapeutic preparation of this invention.

Example 3:

15	Stearic acid	2 wt. %
	Cetyl alcohol	0.8
	Bees wax	1
	Squalane	3
	Olive oil	1
20	Polyoxyethylene(10 moles) monooleate	2
	Glycerin	5
	Ethanol	3
	Carboxyvinyl polymer	0.2
25	Triethanol amine	1.2
	Ubidecarenone	0.1
	Antioxidant	as needed
	Perfume	as needed

0146742
0.3

Butylparaben

Purified water

80.4

The above components were mixed in a manner known
per se in the art into a homogeneous lotion as a
5 therapeutic preparation of this invention.

Example 4:

	Liquid paraffin	5 wt. %
	Vaseline	2
	Cetanol	1
10	Glycerin monostearate	2
	Polyoxyethylene (20 moles) sorbitan monolaurate	1
	Ubidecarenone	0.5
	Dipropylene glycol	5
15	Xanthan gum	1
	Methylparaben	0.2
	Antioxidant	as needed
	Perfume	as needed
	Purified water	82.3

20 The above component were mixed in a manner known
per se in the art into a homogeneous lotion as a
therapeutic preparation of this invention.

Having now fully described the invention, it will be
apparent to one of ordinary skill in the art that many
25 changes and modifications can be made thereto without
departing from the spirit or scope of the invention as set
forth herein.

Claims

0146742

1. In a cutaneously-applicable therapeutic preparation for radiation ulcers, the improvement wherein the preparation contains ubidecarenone as an effective component.

5

2. A preparation as claimed in Claim 1, wherein ubidecarenone is contained in an amount of 0.05 - 5 wt.%.

3. A preparation as claimed in Claim 2, wherein
10 ubidecarenone is contained in an amount of 0.1 - 2.0 wt.%.

4. A preparation as claimed in Claim 1, further comprising an excipient which gives little irritation to the skin.

15

5. A preparation as claimed in Claim 4, wherein the excipient is selected from the group consisting of glycerin, hohoba oil, cetyl alcohol, olive oil, glycerin esters of fatty acids and mixtures thereof.

20

6. A preparation as claimed in Claim 1, wherein the preparation is in the form of a cream.

7. A preparation as claimed in Claim 1, wherein the
25 preparation is in the form of a lotion.

0146742

8. A preparation as claimed in Claim 7, wherein
ubidecarenone is present in combination with cytochrome C
and/or urokinase.



European Patent
Office

EUROPEAN SEARCH REPORT

01 467 42
Application number

EP 84 11 3349

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int Cl 4)
X, Y	GB-A-2 116 426 (SEUREF A.G.) * Column 1, line 33 - column 2, line 73; column 1, lines 11-17; claims *	1-7	A 61 K 31/12 A 61 K 47/00
Y	--- CHEMICAL ABSTRACTS, vol. 98, no. 1, 3rd January 1983, page 213, no. 2293v, Columbus, Ohio, US; A.F. KOZHOKARU et al.: "Modification of radiation damage by coenzyme Q and dimethyl sulfoxide"; & RADIOBIOLOGIYA 1982, 22(5), 693-6 * Abstract *	1-4, 6-7	
Y	--- GB-A-1 317 865 (JULO PAPER CO.) * Page 1, lines 8-19, 70-86; page 3, lines 96-124; page 4, lines 27-31; page 17; claims 1, 12, 39 *	1-7	TECHNICAL FIELDS SEARCHED (Int Cl 4) A 61 K 31/00 A 61 K 47/00

The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 06-03-1985	Examiner BERTE M.J.
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with an th r document of the same category A : technological background O : non-written disclosure P : intermediate d cument		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	